

## **Product datasheet for TA328135S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## 14-3-3 zeta (YWHAZ) Mouse Monoclonal Antibody [Clone ID: 1314CT423.108.153.173.140]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: 1314CT423.108.153.173.140

Applications: WB

Recommended Dilution: WB: 1:1000

**Reactivity:** Human, Mouse, Rat (Predicted: Bovine)

Host: Mouse

Isotype: IgG2b, kappa
Clonality: Monoclonal

**Immunogen:** This YWHAZ antibody is generated from a mouse immunized with a recombinant protein

from human YWHAZ.

**Formulation:** PBS with 0.09% (W/V) sodium azide

**Concentration:** lot specific

**Purification:** This antibody is purified through a protein G column, followed by dialysis against PBS.

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta

Database Link: NP 663723

Entrez Gene 22631 MouseEntrez Gene 25578 RatEntrez Gene 7534 Human

P63104

**Background:** Adapter protein implicated in the regulation of a large spectrum of both general and

specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of

the activity of the binding partner.

Synonyms: 14-3-3-zeta; HEL-S-3; HEL4; KCIP-1; YWHAD


**Protein Pathways:** Cell cycle, Neurotrophin signaling pathway, Oocyte meiosis, Pathogenic Escherichia coli

infection





## **Product images:**



Western blot analysis of lysates from A431, HepG2, mouse NIH/3T3, rat PC-12 and C6 cell line, mouse brain tissue (from left to right), using YWHAZ Antibody (Cat. #[TA328135]). [TA328135] was diluted at 1:1000 at each lane. A goat antimouse IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.