

Product datasheet for TA328138S

i roddet datasiieet for 1A320130.

POLR2C Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:1000

Reactivity: Human, Mouse (Predicted: Bovine)

Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This POLR2C antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 200-228 amino acids from the C-terminal region of human POLR2C.

Formulation: PBS with 0.09% (W/V) sodium azide

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: polymerase (RNA) II subunit C

Database Link: NP 116558

Entrez Gene 20021 MouseEntrez Gene 5432 Human

P19387

Background: DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four

ribonucleoside triphosphates as substrates. Component of RNA polymerase II which synthesizes mRNA precursors and many functional non-coding RNAs. Pol II is the central component of the basal RNA polymerase II transcription machinery. It is composed of mobile elements that move relative to each other. RPB3 is part of the core element with the central large cleft and the clamp element that moves to open and close the cleft (By similarity).

Synonyms: hRPB33; hsRPB3; RPB31

Protein Families: Druggable Genome, Transcription Factors



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Pathways:

Huntington's disease, Metabolic pathways, Purine metabolism, Pyrimidine metabolism, RNA polymerase

Product images:



POLR2C Antibody (C-term) (Cat. #[TA328138]) western blot analysis in Hela, U251 cell line and mouse skeletal muscle tissue lysates (35ug/lane). This demonstrates the POLR2C antibody detected the POLR2C protein (arrow).