

## **Product datasheet for TA328209**

## IL3 Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** E.coli derived Recombinant Human IL-3

**Formulation:** A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Purification:** Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-3.

Anti-Human IL-3 specific antibody was purified by affinity chromatography employing

immobilized hIL-3 matrix.

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Gene Name: interleukin 3

Database Link: NP 000579

Entrez Gene 3562 Human

P08700

Synonyms: IL-3; MCGF; MULTI-CSF

**Note:** Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hIL-3

(1.0 ng/ml), a concentration of 0.38 - 0.57 <=ug/ml of this antibody is required.

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Apoptosis, Asthma, Cytokine-cytokine receptor interaction, Fc epsilon RI signaling pathway,

Hematopoietic cell lineage, Jak-STAT signaling pathway




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



To detect hIL-3 by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human IL-3 ([TA328208]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-3.



To detect hIL-3 by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human IL-3 ([TA328208]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-3.