

## **Product datasheet for TA328257**

## **CD137 (TNFRSF9) Goat Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL

Reactivity: Human
Host: Goat
Isotype: IgG

Clonality: Polyclonal

Immunogen: E.coli derived Recombinant Human 4-1BB Receptor

**Formulation:** A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Purification:** Produced from sera of goats pre-immunized with highly pure (>98%) recombinant human 4-

1BB Receptor. Anti-Human 4-1BB Receptor specific antibody was purified by affinity

chromatography employing immobilized h4-1BB Receptor matrix.

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** tumor necrosis factor receptor superfamily member 9

Database Link: NP 001552

Entrez Gene 3604 Human

Q07011

**Synonyms:** 4-1BB; CD137; CDw137; ILA

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Cytokine-cytokine receptor interaction


**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



To detect h4-1BB Receptor by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human 4-1BB Receptor ([TA328256]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant h4-1BB Receptor.



To detect h4-1BB Receptor by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human 4-1BB Receptor ([TA328256]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant h4-1BB Receptor.