

Product datasheet for **TA328282**

CCL27 Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	WB: 0.1-0.2ug/mL, ELISA: 0.5-2ug/mL, IHC: 0.25ug/mL - 1 mg/mL
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Formulation:	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Purification:	Produced from sera of rabbits immunized with highly pure recombinant Human CTACK. Anti-Human CTACK specific antibody was purified by affinity chromatography employing an immobilized Human CTACK matrix.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	C-C motif chemokine ligand 27
Database Link:	NP_006655 Entrez Gene 10850 Human Q9Y4X3
Synonyms:	ALP; CTACK; CTAK; ESKINE; ILC; PESKY; SCYA27
Protein Families:	Druggable Genome
Protein Pathways:	Chemokine signaling pathway, Cytokine-cytokine receptor interaction




[View online »](#)


Product images:



To detect Human CTACK by sandwich ELISA (using 100ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human CTACK ([TA328281]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human CTACK.



To detect Human CTACK by sandwich ELISA (using 100ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human CTACK ([TA328281]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human CTACK.



To detect Human CTACK by sandwich ELISA (using 100ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human CTACK ([TA328281]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human CTACK.