

## Product datasheet for TA328384

## **MIA2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** ELISA, IHC, WB

Recommended Dilution: WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL, IHC: 0.25ug/mL - 1mg/mL

Reactivity: Rabbit Host: Isotype: lgG

Clonality: Polyclonal

Immunogen: E.coli derived Recombinant Human MIA-2

Formulation: A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Purification:** Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hMIA-2

(human MIA-2). Anti-Human MIA-2 specific antibody was purified by affinity chromatography

employing immobilized hMIA-2 matrix.

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Gene Name: melanoma inhibitory activity 2

Database Link: NP 473365

Entrez Gene 4253 HumanEntrez Gene 117153 Human

Synonyms: FLI22404

**Protein Families:** Transmembrane




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




## **Product images:**



To detect hMIA-2 by sandwich ELISA (using 100ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human MIA-2 ([TA328383]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIA-2.



To detect hMIA-2 by sandwich ELISA (using 100ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human MIA-2 ([TA328383]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIA-2.



To detect hMIA-2 by sandwich ELISA (using 100ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Human MIA-2 ([TA328383]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIA-2.