

## Product datasheet for **TA328426**


### **RANKL (TNFSF11) Rabbit Polyclonal Antibody**

#### **Product data:**

<b>Product Type:</b>	Primary Antibodies
<b>Applications:</b>	WB
<b>Recommended Dilution:</b>	WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL
<b>Reactivity:</b>	Human
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Clonality:</b>	Polyclonal
<b>Immunogen:</b>	E.coli derived Recombinant Human sRANK Ligand
<b>Formulation:</b>	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
<b>Purification:</b>	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hsRANKL. Anti-Human sRANKL specific antibody was purified by affinity chromatography and then biotinylated.
<b>Conjugation:</b>	Biotin
<b>Storage:</b>	Store at -20°C as received.
<b>Stability:</b>	Stable for 12 months from date of receipt.
<b>Gene Name:</b>	tumor necrosis factor superfamily member 11
<b>Database Link:</b>	<a href="#">NP_003692</a> <a href="#">Entrez Gene 8600 Human</a> <a href="#">O14788</a>
<b>Synonyms:</b>	CD254; hRANKL2; ODF; OPGL; OPTB2; RANKL; sOdf; TRANCE
<b>Protein Families:</b>	Druggable Genome, Transmembrane
<b>Protein Pathways:</b>	Cytokine-cytokine receptor interaction



[View online »](#)

**Product images:**

To detect hsRANKL by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 ug/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hsRANKL is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.