

Product datasheet for **TA328442**


TACI (TNFRSF13B) Goat Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	ELISA
Recommended Dilution:	WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL
Reactivity:	Human
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	E.coli derived Recombinant Human TACI (TNFRSF13B)
Formulation:	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Purification:	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hTACI. Anti-Human TACI specific antibody was purified by affinity chromatography and then biotinylated.
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	tumor necrosis factor receptor superfamily member 13B
Database Link:	NP_036584 Entrez Gene 23495 Human O14836
Synonyms:	CD267; CVID; CVID2; IGAD2; RYZN; TACI; TNFRSF14B
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, Primary immunodeficiency



[View online »](#)

Product images:


To detect hTAC1 by direct ELISA (using 100 ul/well antibody solution) a concentration of 0.25-1.0 ug/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2-0.4 ng/well of recombinant hTAC1.



To detect hTAC1 by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.25-1.0 ug/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with Polyclonal Anti-Human TAC1 ([TA328443]) as a capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hTAC1.