

## Product datasheet for **TA328457**


### **TNFRSF1A Rabbit Polyclonal Antibody**

#### **Product data:**


|                              |   |
|------------------------------|---|
| <b>Product Type:</b>         | Primary Antibodies  |
| <b>Applications:</b>         | ELISA   |
| <b>Recommended Dilution:</b> | WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL  |
| <b>Reactivity:</b>           | Human   |
| <b>Host:</b>                 | Rabbit  |
| <b>Isotype:</b>              | IgG   |
| <b>Clonality:</b>            | Polyclonal  |
| <b>Immunogen:</b>            | E.coli derived Recombinant Human sTNF Receptor Type I   |
| <b>Formulation:</b>          | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.  |
| <b>Purification:</b>         | Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hsTNF-Receptor I. Anti-Human sTNF-Receptor I specific antibody was purified by affinity chromatography and then biotinylated. |
| <b>Conjugation:</b>          | Biotin  |
| <b>Storage:</b>              | Store at -20°C as received.   |
| <b>Stability:</b>            | Stable for 12 months from date of receipt.  |
| <b>Gene Name:</b>            | tumor necrosis factor receptor superfamily member 1A  |
| <b>Database Link:</b>        | <a href="#">NP_001056</a><br><a href="#">Entrez Gene 7132 Human</a><br><a href="#">P19438</a>   |
| <b>Synonyms:</b>             | CD120a; FPF; MS5; p55; p55-R; p60; TBP1; TNF-R; TNF-R-I; TNF-R55; TNFAR; TNFR1; TNFR1-d2; TNFR55  |
| <b>Protein Families:</b>     | Druggable Genome, Secreted Protein, Transcription Factors, Transmembrane  |
| <b>Protein Pathways:</b>     | Adipocytokine signaling pathway, Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Apoptosis, Cytokine-cytokine receptor interaction, MAPK signaling pathway  |



[View online »](#)

**Product images:**


To detect hsTNF-Receptor I by direct ELISA (using 100 ul/well antibody solution) a concentration of 0.25-1.0 ug/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2-0.4 ng/well of recombinant hsTNF-Receptor I.



To detect hsTNF-Receptor I by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.25-1.0 ug/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with Polyclonal Anti-Human sTNF-Receptor I ([TA328458]) as a capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant hsTNF-Receptor I.