

## **Product datasheet for TA328534**

## **Ccl12 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL

Reactivity: Murine
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: E.coli derived Recombinant Murine MCP-5 (CCL12)

**Formulation:** A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Purification:** Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mMCP-5.

Anti-Murine MCP-5 specific antibody was purified by affinity chromatography and then

biotinylated.

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** chemokine (C-C motif) ligand 12

Database Link: NP 035461

Entrez Gene 20293 Mouse

Q62401

Synonyms: MCP-5; Scya12




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




## **Product images:**



To detect mMCP-5 by direct ELISA (using 100 ul/well antibody solution) a concentration of 0.25-1.0 ug/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2-0.4 ng/well of recombinant mMCP-5.



To detect mMCP-5 by direct ELISA (using 100 ul/well antibody solution) a concentration of 0.25-1.0 ug/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2-0.4 ng/well of recombinant mMCP-5.



To detect mMCP-5 by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.25-1.0 ug/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with Polyclonal Anti-Murine MCP-5 ([TA328535]) as a capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant mMCP-5.