

Product datasheet for **TA328547**


Tnfsf11 Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL
Reactivity:	Murine
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	E.coli derived Recombinant Murine sRANK Ligand
Formulation:	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Purification:	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant msRANKL. Anti-Murine sRANKL specific antibody was purified by affinity chromatography employing immobilized msRANKL matrix.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	tumor necrosis factor (ligand) superfamily, member 11
Database Link:	NP_035743 Entrez Gene 21943 Mouse O35235
Synonyms:	CD254; hRANKL2; ODF; OPGL; OPTB2; OTTHUMP00000178585; RANKL; sOdf; TRANCE
Note:	Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of msRANKL (50.0 ng/ml), a concentration of 0.05 ug/ml of this antibody is required.


[View online »](#)

Product images:



To detect msRANKL by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Murine sRANKL ([TA328546]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant msRANKL.



To detect msRANKL by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Murine sRANKL ([TA328546]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant msRANKL.