

## **Product datasheet for TA328555**

## **Retn Goat Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL

Reactivity: Murine
Host: Goat
Isotype: IgG

Clonality: Polyclonal

Immunogen: E.coli derived Recombinant Murine Resistin

**Formulation:** A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Purification:** Produced from sera of goats pre-immunized with highly pure (>98%) recombinant mResistin.

Anti-Murine Resistin specific antibody was purified by affinity chromatography employing

immobilized mResistin matrix.

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** resistin

Database Link: NP 075360

Entrez Gene 57264 Mouse

Q99P87

Synonyms: ADSF; FIZZ3; HXCP1; MGC126603; MGC126609; resistin; RETN1; RSTN; XCP1




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



To detect mResistin by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Murine Resistin ([TA328554]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mResistin.



To detect mResistin by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Murine Resistin ([TA328554]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mResistin.