

Product datasheet for **TA328561**

Vegfa Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	WB: 0.1-0.2ug/mL, ELISA: 0.25-2ug/mL, IHC: 0.25ug/mL - 1mg/mL
Reactivity:	Murine
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	E.coli derived Recombinant Murine VEGF
Formulation:	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Purification:	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mVEGF. Anti-Murine VEGF specific antibody was purified by affinity chromatography employing immobilized mVEGF matrix.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	vascular endothelial growth factor A
Database Link:	NP_033531 Entrez Gene 22339 Mouse Q00731
Synonyms:	MGC70609; MVCD1; VEGF; VEGF-A; VPF
Note:	Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of mVEGF (10.00 ng/ml), a concentration of 0.08-0.12 <=ug/ml of this antibody is required.




[View online »](#)


Product images:



To detect mVEGF by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Murine VEGF ([TA328560]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mVEGF.



To detect mVEGF by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Murine VEGF ([TA328560]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mVEGF.



To detect mVEGF by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biotinylated Anti-Murine VEGF ([TA328560]) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mVEGF.