

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA328610

NPY1R Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	WB: 1:200-1:2000; IHC: 1:100-1:3000
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide (C)RLKRRNNMMDKMRDNK, corresponding to amino acid residues 237-252 of human NPY1R. 3rd intracellular loop.
Formulation:	Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to CoA along with shipment for actual concentration). Buffer before lyophilization: Phosphate buffered saline (PBS), pH 7.4, 1% BSA, 0.05% NaN3.
Reconstitution Method:	Add 50 ul double distilled water (DDW) to the lyophilized powder.
Purification:	Affinity purified on immobilized antigen.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	neuropeptide Y receptor Y1
Database Link:	<u>NP 000900</u>
	<u>Entrez Gene 29358 RatEntrez Gene 4886 Human</u> <u>P25929</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE NPY1R Rabbit Polyclonal Antibody – TA328610

Background:

Neuropeptide Y (NPY) is a 36-amino acid peptide neurotransmitter in the central (CNS) and peripheral nervous systems. It belongs to the NPY family, together with the peptide YY (PYY) and the pancreatic polypeptide (PP). NPY is involved in regulation of a broad range of homeostatic functions such as eating behavior, hypertension, and reproduction. NPY exerts its effects via five receptor subtypes known as the neuropeptide Y receptors (NPYR): Y1 (NPY1R), Y2 (NPY2R), Y4 (NPY4R), Y5 (NPY5R) and Y6 (NPY6R). Four of these, Y1, Y2, Y4, and Y5, are expressed in humans. The NPY receptors belong to the G protein-coupled receptor superfamily whose members share a common structure of seven putative transmembrane domains, an extracellular amino terminus, and a cytoplasmic carboxyl terminus. Activation of these receptors by NPY produces effects on [Ca2+], adenylate cyclase, and a number of ion channels.Based on the high frequency and density of NPY receptors in steroid hormoneproducing tumors, the NPY receptors might be of potential use in tumor management. NPY1R is considered to be a postsynaptic receptor. In the CNS, NPY1Rs are concentrated in the cortical areas of the brain, cerebral cortex, olfactory nucleus, hippocampus, hypothalamic nuclei, and thalamic nuclei. In addition, this receptor has a bimodal role in the immune system, serving as a strong negative regulator on T cells as well as a key activator of antigenpresenting cell function.

Synonyms:	NPY1-R; NPYR
Protein Families:	Druggable Genome, GPCR, Transmembrane
Protein Pathways:	Neuroactive ligand-receptor interaction

Product images:



Western blot analysis of Jurkat (lanes 1 and 4), K562 (lanes 2 and 5), and RBL (lanes 3 and 6) cell lysates and rat brain lysates (lanes 7 and 8): 1, 2, 3, 7. Anti-Neuropeptide Y1 Receptor antibody, (1:200). 4, 5, 6, 8. Anti-Neuropeptide Y1 Receptor antibody preincubated with the control peptide antigen.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunocytochemical staining of a primary culture of rat dorsal root ganglion (DRG) neurons. A, D. A paraformaldehyde-fixed and permeabilized DRG primary culture was stained with Anti-Neuropeptide Y1 Receptor antibody, (1:100), followed by Alexa-555-conjugated goatanti-rabbit secondary antibody. B, E. Nuclear fluorescence staining of cells using the membrane-permeable DNA dye Hoechst 33342. C. Merged images of A and B. F. Merged images of D and E. Magnification: A-C: x20 E-F: x100

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US