

Product datasheet for TA328621

KCNK3 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:200-1:2000; IHC: 1:100-1:3000
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide (C)EDEKRDAEHRALLTRNGQ, corresponding to amino acid residues 252-269 of human K2P3.1 (TASK-1). Intracellular, C-terminal part.
Formulation:	Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to CoA along with shipment for actual concentration). Buffer before lyophilization: Phosphate buffered saline (PBS), pH 7.4, 1% BSA, 5% sucrose, 0.025% NaN3.
Reconstitution Method:	Add 50 ul double distilled water (DDW) to the lyophilized powder.
Purification:	Affinity purified on immobilized antigen.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	potassium two pore domain channel subfamily K member 3
Database Link:	<u>NP 002237</u> <u>Entrez Gene 29553 Rat</u> <u>O14649</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE KCNK3 Rabbit Polyclonal Antibody – TA328621

Background: K2P3.1 (also named TWIK-Related Acid Sensitive K+ channel, TASK-1 or KCNK3) is a member of the 2-pore (2P) domain K+ channels family that includes at least 16 members. These channels show little time or voltage dependence and are considered to be a??leaka? or a?? backgroundâ? K+ channels, thereby generating background currents which help set the membrane resting potential and cell excitation. The K2P channels have a signature topology that includes four transmembrane domains and two pore domains with intracellular N- and C termini. The functional channel is believed to be a dimer. K2P channels are regulated by diverse physical and chemical stimuli including temperature, changes in intracellular pH, mechanical stretch, inhalation anesthetics, etc. The channels can then be subclassified based in their specific activators. K2P3.1 is extremely sensitive to variations in external pH, a drop in pH from 7.7 to 6.7 will be enough to close most K2P3.1 channels. K2P3.1 channel distribution is relatively wide with high expression in the brain and heart as well as in the pancreas, kidney and lung. K2P3.1 can exist in the brain either as a homodimer or as a heterodimer with the closely related channel K2P9.1 (TASK-3). During periods of high neuronal activity a transient increase in extracellular pH has been recorded. This pH increase may stimulate K2P3.1 channel activity and hence enhance outward K+ currents. This in turn will decrease excitability in these neurons and work as a feedback mechanism.

Synonyms: K2p3.1; OAT1; PPH4; TASK; TASK-1; TBAK1

Protein Families: Druggable Genome, Ion Channels: Potassium, Transmembrane

Product images:



Western blot analysis of rat brain (1, 3) and pancreas (2, 4) membranes: 1, 2. Anti-K2P3.1 (TASK-1) antibody, (1:200). 3, 4. Anti-K2P3.1 (TASK-1) antibody, preincubated with the control peptide

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US