

Product datasheet for **TA328647**

SAP102 (DLG3) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:200-1:2000
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	GST fusion protein of the sequence KSTPKLNGSGPSWW PECTCTNRDWYEQVNGSD, corresponding to amino acid residues 93-124 of human SAP102 (MW: 30 kDa.). N-terminal part.
Formulation:	Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to CoA along with shipment for actual concentration). Buffer before lyophilization: Phosphate buffered saline (PBS), pH 7.4, 1% BSA, 0.025% NaN ₃ .
Reconstitution Method:	Add 50 ul double distilled water (DDW) to the lyophilized powder.
Purification:	The serum was depleted of anti-GST antibodies by affinity chromatography on immobilized GST and then the IgG fraction was purified on immobilized Protein A.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	discs large MAGUK scaffold protein 3
Database Link:	NP_066943 Entrez Gene 58948 Rat Q92796




[View online »](#)

Background:

SAP102 (also known as DLG3) is a PDZ containing domain protein that is also a member of the membrane-associated guanylate kinase (MAGUK) family of multi-domain adaptor proteins. PDZ domains are conserved protein domains of about 90 amino acids involved in protein-protein recognition, protein targeting and assembly of multi-protein complexes. The name PDZ derives from the first three proteins in which these domains were identified: PSD-95 (a 95 kDa protein involved in signaling at the post-synaptic density), DLG (the *Drosophila melanogaster* Discs Large protein) and ZO-1 (the zonula occludens 1 protein involved in maintenance of epithelial polarity). MAGUKs are scaffolding proteins that comprise several modular protein binding motifs including one or more PDZ domains, a Src homology 3 (SH3) domain, and a catalytically inactive guanylate kinase-like domain. The multidomain nature of PDZ-containing proteins enables them to interact with multiple binding partners and hence organize larger signaling protein complexes. SAP102 has been shown to participate in the postsynaptic density, a dedicated structure formed in postsynaptic nerve terminals that includes a specialized assembly of ion channels, receptors and signaling molecules that are involved in information processing and the modulation of synaptic plasticity. Moreover, mutations in the SAP102 gene have been linked with a form of mental retardation.

Synonyms:

MRX; MRX90; NEDLG; PPP1R82; SAP102; XLMR

Product images:


Western blot analysis of rat brain membranes: 1. Anti-SAP 102 antibody, (1:400). 2. Anti-SAP 102 antibody, preincubated with the control fusion protein antigen.