

Product datasheet for TA328772

Trpv4 Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	WB: 1:200-1:2000; IHC: 1:100-1:3000
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide (C)EDQSN(S)TVPSYPA(S)RD, corresponding to amino acid residues 647-662 of rat TRPV4. 3rd extracellular loop.
Formulation:	Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to CoA along with shipment for actual concentration). Buffer before lyophilization: Phosphate buffered saline (PBS), pH 7.4, 1% BSA, 0.05% NaN ₃ .
Reconstitution Method:	Add 50 ul double distilled water (DDW) to the lyophilized powder.
Purification:	Affinity purified on immobilized antigen.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	transient receptor potential cation channel, subfamily V, member 4
Database Link:	NP_076460 Entrez Gene 59341 Human Entrez Gene 66026 Rat Q9ERZ8
Background:	TRP channels are a large family (about 28 genes) of plasma membrane, non-selective cationic channels that are either specifically or ubiquitously expressed in excitable and non-excitable cells. According to IUPHAR the TRP family comprises of three main subfamilies on the basis of sequence homology; TRPC, TRPM and TRPV (to date, three extra subfamilies are considered to belong to the TRP family; the TRPA, TRPML, and TRPP). The TRPV subfamily consists of six members, TRPV1-6. TRPV4 (also named OTRPC4) is activated under hypotonic conditions and serves as an osmoreceptor. TRPV4 is expressed in brain, liver, kidney, heart, testis and salivary gland.


[View online »](#)


Synonyms: OTRPC4; TRP12; VR-OAC; VRL-2; VRL2; VROAC

Note: This antibody was tested in live cell imaging. Please see IF/ICC data for detail.


Product images:




Western blot analysis of rat brain lysate: 1. Anti-TRPV4 (extracellular) antibody, (1:200). 2. Anti-TRPV4 (extracellular) antibody, preincubated with the control peptide antigen.



Expression of TRPV4 in rat DRG.
Immunohistochemical staining of rat dorsal root ganglion (DRG) frozen sections using Anti-TRPV4 (extracellular) antibody. A. TRPV4 labeling appears in neuronal cell bodies. B. Nuclear staining using DAPI as counter stain. C. merged images of A and B.



Expression of TRPV4 in human U-87 cells.
Immunocytochemical staining of live intact human brain glioblastoma U-87 cells. Extracellular staining of cells using Anti-TRPV4 (extracellular) antibody, (1:25), (red) followed by goat anti-rabbit-AlexaFluor-594 secondary antibody. DAPI was used as the counterstaining (blue).



Expression of TRPV4 in human U-87 cells.
Immunocytochemical staining of live intact human brain glioblastoma U-87 cells. A. Extracellular staining of cells with Anti-TRPV4 (extracellular) antibody, (1:25) followed by goat anti-rabbit-AlexaFluor-594 secondary antibody. B. Merged images of live cells with A.