

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA328808

Ffar1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:200-1:2000
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide, (C)NVASFINPDLEGSWRK corresponding to amino acid residues 244-259 of rat free fatty acid receptor 1.
Formulation:	Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to CoA along with shipment for actual concentration). Buffer before lyophilization: Phosphate buffered saline (PBS), pH 7.4, 1% BSA, 0.05% NaN3.
Reconstitution Method:	Add 50 ul double distilled water (DDW) to the lyophilized powder.
Purification:	Affinity purified on immobilized antigen.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	free fatty acid receptor 1
Database Link:	<u>NP 695216</u> <u>Entrez Gene 266607 Rat</u> <u>Q8K3T4</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE Ffar1 Rabbit Polyclonal Antibody – TA328808

Background:

Free fatty acids (FFAs) were for long believed to influence cellular metabolism. Hoever, the discovery that FFAs actually mediate their effects through one of three G-protein coupled receptors (GPCRs) - Free Fatty Acid Receptors 1-3 (FFAR1-3), has initiated a number of studies in order to assess their implication in health and disease. Like all other GPCRs, FFAR1-3 have seven transmembrane domains, an extracellular N-terminal tail and an intracellular Cterminus. FFAR1 is activated by either medium or long, saturated or unsaturated fatty acids, whereas FFAR2 and FFAR3 are activated by short chain fatty acids. FFAR1 is coupled to Gg and leads to Ca2+ mobilization upon activation. FFAR2 couples to both pertussis-sensitive and insensitive G-proteins and also leads to increased intracellular Ca2+ levels. FFAR3 coupling to G-protein induces an increase in cAMP. Distribution studies of all three FFA Receptors indicate that FFAR1 mRNA is guite elevated in the pancreas. FFAR1 is also detected in the liver, heart, skeletal muscle, and brain; although this broad localization pattern is supported only by one study. FFAR2 is expressed in many tissues like liver and colon. FFAR3 is expressed in many tissues but shows highest expression in adipose tissue. Relatively high expression of the receptor could be detected in pancreas, spleen, lymph nodes and bone marrow. As mentioned above, the finding that FFAsâ?? actions are mediated by GPCRs, prompted many studies in order to elucidate their roles in healthy and disease states. Such studies stipulate that FFAR1 may be involved in obesity and type 2 diabetes since circulating free fatty acid levels in the plasma are significantly elevated. On the other hand, FFAR2 and FFAR3 could be potential targets for treating irritable bowel disease and appetite control.

Synonyms:

FFA1R; GPCR40; GPR40

Product images:



Western blot analysis of rat pancreas lysate: 1. Anti-Free Fatty Acid Receptor 1 (extracellular) antibody, (1:200). 2. Anti-Free Fatty Acid Receptor 1 (extracellular) antibody, preincubated with the control peptide antigen.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US