

Product datasheet for **TA328826**

Grm5 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	WB: 1:200-1:2000; IHC: 1:100-1:3000
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide EGFAQENSKYNKTC, corresponding to amino acids 367-380 of rat mGluR5. Extracellular, N-terminus.
Formulation:	Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to CoA along with shipment for actual concentration). Buffer before lyophilization: Phosphate buffered saline (PBS), pH 7.4, 1% BSA, 0.025% NaN ₃ .
Reconstitution Method:	Add 50 ul double distilled water (DDW) to the lyophilized powder.
Purification:	Affinity purified on immobilized antigen.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	glutamate metabotropic receptor 5
Database Link:	NP_058708 Entrez Gene 108071 Mouse Entrez Gene 24418 Rat P31424



[View online »](#)

Background:


L-Glutamate, the major excitatory neurotransmitter in the central nervous system, operates through several receptors that are categorized as ionotropic (ligand-gated cation channels) or metabotropic (G-protein coupled receptors). The metabotropic glutamate receptors family includes eight members (mGluR1-8) that have been divided into three groups based on their sequence homology, pharmacology and signal transduction. Group I of the metabotropic glutamate receptors includes the mGluR1 and mGluR5 receptors. The receptors present the typical G-protein coupled receptor (GPCR) signature topology: seven transmembrane domains with a large extracellular N-terminus domain and an intracellular C-terminus one. The N-terminus domain of group I receptors contains the glutamate binding site while the cytoplasmic C-terminus domain has an important role in the regulation of receptor activity through interactions with other proteins such as the Homer adaptor proteins. mGluR1 and mGluR5 receptors signal through Gq/G11 that activates phospholipase C and ultimately produces an increase in inositol trisphosphate and cytosolic Ca²⁺. More downstream signaling pathways include activation of PKC and modulation of Ca²⁺ and K⁺ ion channels. Activation of signaling pathways independent of G-proteins has also been reported. mGluR5 is predominantly expressed in nervous tissue although expression in several non-neural cell types has also been described. In the brain it is highly expressed in the cortex, basal ganglia and hippocampus. The mGluR5 receptor is involved in several physiological processes such as neuronal development, induction of long-term potentiation (LTP) and depression (LTD) as well as in pathological disorders such as brain trauma, chronic pain, Parkinson's and Huntington's disease.

Synonyms:


GPRC1E; mGlu5; MGLUR5; OTTHUMP00000165953; OTTHUMP00000165957

Note:


This antibody was tested in live cell imaging. Please see IF/ICC data for detail.

Product images:


Western blot analysis of rat brain membranes: 1. Anti-mGluR5 (extracellular) antibody, (1:500). 2. Anti-mGluR5 (extracellular) antibody, preincubated with the control peptide antigen.



IHC staining of perfusion-fixed frozen rat cerebellum sections using Anti-mGluR5 (extracellular) antibody, (1:50). mGluR5 (red) was detected in cerebellar Purkinje cells (vertical arrows) and in the molecular layer (horizontal arrows). Staining with mouse anti-parvalbumin (green) revealed co-localization in Purkinje but not in the molecular layer. Little staining of mGluR5 was detected in the granule layer. DAPI counterstain is used to visualize nuclei of all cells (blue).



Expression of mGluR5 in rat cerebellum and hippocampus. Immunohistochemical staining of perfusion-fixed frozen rat hippocampus sections using Anti-mGluR5 (extracellular) antibody, (1:50). mGluR5 (red) was detected in CA3 cells (arrows). Staining with mouse anti-parvalbumin (green) revealed co-localization in pyramidal layer. DAPI counterstain was used to visualize nuclei of all cells (blue).



Expression of mGluR5 in rat GH3 pituitary cells. Immunocytochemical staining of live intact rat GH3 pituitary cells using Anti-mGluR5 (extracellular) antibody, (1:100), followed by goat-anti-rabbit-AlexaFluor-555 secondary antibody.