

Product datasheet for **TA328840**

Slc1a6 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:200-1:2000
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide (C)TRTIVRTDNGSE, corresponding to amino acid residues 205- 216 of rat EAAT4. 2nd extracellular loop.
Formulation:	Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to CoA along with shipment for actual concentration). Buffer before lyophilization: Phosphate buffered saline (PBS), pH 7.4, 1% BSA, 0.05% NaN ₃ .
Reconstitution Method:	Add 50 ul double distilled water (DDW) to the lyophilized powder.
Purification:	Affinity purified on immobilized antigen.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	solute carrier family 1 member 6
Database Link:	NP_114454 Entrez Gene 20513 Mouse Entrez Gene 84012 Rat O35921




[View online »](#)

Background:

L-Glutamate (Glu) is an abundant amino acid that functions as the major excitatory neurotransmitter in the central nervous system. However, excess of Glu in the extracellular synaptic milieu leads to neuronal cell death by a process known as excitotoxicity. The extracellular levels of Glu are regulated by a family of high affinity plasma membrane transporters called excitatory amino acid transporters (EAATs) which are responsible for the re-uptake of Glu into the cells. The EAAT family includes five members (EAAT1-EAAT5) that are members of the solute carrier family 1 (SLC1) of Na⁺-dependent transporters that also includes the neutral amino acid transporters ASCT1 and ASCT2. The Glu transporters present an unusual topology of eight transmembrane domains with two re-entrant loops and intracellular N- and C- termini. The transporter is likely assembled as a trimer where each monomer is a functional unit capable of binding the Glu substrate. The transport of Glu into the cells by the EAAT transporters is coupled to the Na⁺ and K⁺ electrochemical gradient as a driving force. Hence, the uptake of Glu is dependent on the co-transport of three Na⁺ and one H⁺ ions, and the counter transport of one K⁺ ion. In addition, to the well documented Glu uptake, the EAAT transporters show a Glu-independent Cl⁻ conductance. The physiological significance of the Cl⁻ current through the EAATs is currently unknown. EAAT2 as well as EAAT1 are expressed predominantly in glia cells, while EAAT3 (Excitatory Amino Acid Transporter 3), EAAT4 and EAAT5 are mostly expressed in neurons. As mentioned, EAAT transporters represent the only (significant) mechanism for removal of glutamate from the extracellular fluid and hence are essential for the long-term maintenance of low and non-toxic concentrations of glutamate and the preservation of normal excitatory synaptic transmission. In addition to Glu uptake, the glutamate transporters provide glutamate for the synthesis of g-Aminobutyric acid (GABA), glutathione and protein, suggesting an interactive role between EAATs and cellular metabolism. Dysregulation of EAAT activities has been implicated in several neurodegenerative disorders such as Alzheimer's disease, traumatic brain injury, epilepsy and schizophrenia, suggesting that EAATs can be a useful target for the treatment of these conditions.

Synonyms:

EAAT4; MGC33092; MGC43671

Product images:

Western blot analysis of mouse (lanes 1 and 3) and rat (lanes 2 and 4) brain lysate: 1-2. Anti-Excitatory Amino Acid Transporter 4 (extracellular) antibody, (1:200). 3-4. Anti-Excitatory Amino Acid Transporter 4 (extracellular) antibody, preincubated with the control peptide antigen.