

Product datasheet for **TA328893**

Stx3 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:200-1:2000
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide KDRLEQLKAKQLTQD(C), corresponding to amino acid residues 2-16 of rat syntaxin 3. Cytoplasmic, N-terminus.
Formulation:	Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to CoA along with shipment for actual concentration). Buffer before lyophilization: Phosphate buffered saline (PBS), pH 7.4, 1% BSA, 5% sucrose, 0.025% NaN ₃ .
Reconstitution Method:	Add 50 ul double distilled water (DDW) to the lyophilized powder.
Purification:	Affinity purified on immobilized antigen.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	syntaxin 3
Database Link:	NP_112386 Entrez Gene 81802 Rat Q08849




[View online »](#)

Background:

Syntaxin 3 is a member of the soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) protein superfamily. The family includes 36 members in humans and is characterized by the SNARE motif, an evolutionarily conserved stretch of 60–70 amino acids that are arranged in heptad repeats. SNARE proteins are involved in exocytosis and intracellular vesicle trafficking and are essential for cell growth, hormone secretion and neurotransmission, processes that require rapid, targeted, and regulated membrane fusion. SNAREs can be roughly divided into vesicular (v-SNAREs) and target (t-SNAREs) based on their distribution on the transport vesicle or target membrane respectively. Thus, assembly of cognate v-/t-SNAREs between two opposing membranes generates trans-SNARE complexes, which bring the lipid bilayers in close proximity and drive membrane fusion. Syntaxin 3, like most SNAREs, is a type IV membrane protein with a relatively large N-terminus containing the SNARE motif located in the cytoplasmic side and a transmembrane domain located close to the C-terminus that functions as an anchor. Syntaxin 3 functions as a plasma membrane protein t-SNARE that is involved in epithelial cell polarization. Indeed, Syntaxin 3 is specifically localized in the apical membrane of polarized epithelial cells where it likely has a role in the polarized trafficking of newly synthesized proteins.

Synonyms:

STX3A

Product images:

Western blot analysis of rat brain (lanes 1 and 3) and kidney (lanes 2 and 4) membranes: 1, 2. Anti-Syntaxin 3 antibody, (1:200). 3, 4. Anti-Syntaxin 3 antibody, preincubated with the control peptide antigen.