

Product datasheet for TA328949

Product datasireet for TAS2694

Kcnq2 Rabbit Polyclonal Antibody

Product data:

Clonality:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB: 1:200-1:2000; IHC: 1:100-1:3000

Polyclonal

Reactivity: Human, Mouse, Rat

Host: Rabbit

Immunogen: Peptide (C)RGPTITDKDRTKGPAE, corresponding to amino acid residues 578-593 of rat KCNQ2

. Intracellular, C-terminus.

Formulation: Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to

CoA along with shipment for actual concentration). Buffer before lyophilization: phosphate

buffered saline (PBS), pH 7.4, 1% BSA, 0.05% NaN3.

Reconstitution Method: Add 50 ul double distilled water (DDW) to the lyophilized powder.

Purification: Affinity purified on immobilized antigen.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: potassium voltage-gated channel subfamily Q member 2

Database Link: NP 579856

Entrez Gene 3785 HumanEntrez Gene 16536 MouseEntrez Gene 170848 Rat

088943



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The KCNQ family of voltage-gated K+ channels includes 5 known members: KCNQ1 to KCNQ5. Structurally, the KCNQ family belongs to the six transmembrane domain category of K+ channels. KCNQ family members can form either homomultimeric or heteromultimeric channels with different functional consequences. For example KCNQ2 and KCNQ3 heteromultimers give rise to a much larger channel current than when either protein is expressed alone, probably due to enhanced plasma membrane expression of the combined channel. Indeed, KCNQ2/KCNQ3 heteromultimers are believed to be the molecular correlates of the so-called M current. This current is a K+ neuronal current that is strongly inhibited by the activation of the M1 subtype of the muscarinic acetylcholine receptor. Mutations in either KCNQ2 or KCNQ3 are associated with a form of epilepsy known as benign familial neonatal convulsions (BNFC).

Synonyms:

BFNC; EBN; EBN1; ENB1; HNSPC; KCNA11; KV7.2; KVEBN1

Product images:



Western blot analysis of Kv7.2 transfected HEK-293 cells: 1. Anti-Kv7.2 (KCNQ2) antibody, (1:200). 2. Anti-Kv7.2 (KCNQ2) antibody, preincubated with the control peptide antigen.