

# Product datasheet for TA328951

# **HCN4 Rabbit Polyclonal Antibody**

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies	
Applications:	IHC, WB	
Recommended Dilution:	WB: 1:200-1:2000; IHC: 1:100-1:3000	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Immunogen:	GST fusion protein with sequence HGHLHDSAEERRLIAEGDASPG EDRTPPGLAAEPERP, corresponding to amino acid residues 119-155 of human HCN4 , (MW: 31 kDa.).? ?Â Intracellular, N-terminus.	
Formulation:	Lyophilized. Concentration before lyophilization ~0.8mg/ml (lot dependent, please refer to CoA along with shipment for actual concentration). Buffer before lyophilization: phosphate buffered saline (PBS), pH 7.4, 1% BSA, 0.05% NaN3.	
<b>Reconstitution Method:</b>	Add 50 ul double distilled water (DDW) to the lyophilized powder.	
Purification:	The serum was depleted of anti-GST antibodies by affinity chromatography on immobilized GST, and then anti-HCN4 antibody was affinity purified on immobilized HCN4-GST.	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Gene Name:	hyperpolarization activated cyclic nucleotide gated potassium channel 4	
Database Link:	<u>NP_005468</u> Entrez Gene 59266 RatEntrez Gene 330953 MouseEntrez Gene 10021 Human Q9Y3Q4	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** HCN4 Rabbit Polyclonal Antibody – TA328951

Background: Hyperpolarization-activated cation currents (IHCC) appear in the heart and the brain and have a crucial role in controlling electrical pacemaker activity, contributing to biological processes such as heartbeat, sleep-wake cycle and synaptic plasticity. The IHCC currents are generated by the Hyperpolarization-activated cyclic nucleotide-gated channel family (HCN), which is comprised of four homologous members, HCN1-4. Each HCN subunit consists of six transmembrane domains (TM), a pore region between TM5-TM6 and a binding domain for cyclic nucleotides (CNBD) in the cytoplasmic C-terminus. The HCN subunits can form functional homomers and can also co-assemble into functional heteromers. The channels are closely related to each other and share a homology of about 60%. However, their similarity decreases in the cytoplasmic N- and C-termini. The HCN1-4 channels mainly differ from each other in their speed of activation and the extent to which they are modulated by cAMP. HCN1, weakly affected by cAMP, is the fastest channel, followed by HCN2, HCN3 and HCN4. HCN4 is highly expressed in a restricted manner in adult sinoatrial (SA) node, constituting a good molecular marker for the adult cardiac pacemaker and might serve as a unique marker of the developing SA node . mRNA expression of HCN4 is most abundant in medial habenula and anterior and principal relay nuclei of the thalamus.

Synonyms:	SSS2
Protein Families:	Druggable Genome, Ion Channels: Cyclic nucleotide gated, Transmembrane

### **Product images:**

1 2	
	- 250
	- 130
	- 95
	- 72
-	- 55
	- 36

Western blot analysis of rat brain membranes: 1. Anti-HCN4 antibody, (1:200). 2. Anti-HCN4 antibody, preincubated with the control peptide antigen.



Expression of HCN4 in mouse thalamus. Immunohistochemical staining mouse thalamus using Anti-HCN4 antibody. A. HCN4 (red) appears in the neuropil of the lateral nucleus (LN). B. Staining of reticular nucleus (RN) with mouse anti-Parvalbumin. C. Confocal merge of HCN4 and Parvalbumin images demonstrates separate localization of these proteins in the thalamus.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US