

Product datasheet for TA329080

ATP5PB Goat Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Primary Antibodies |
|-----------------------|--|
| Applications: | WB |
| Recommended Dilution: | WB: 0.03-0.1ug/ml, ELISA: 1:128,000 |
| Reactivity: | Human |
| Host: | Goat |
| lsotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Internal region (SIQHIQNAIDTE) |
| Formulation: | Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. |
| Concentration: | lot specific |
| Purification: | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Gene Name: | ATP synthase, H+ transporting, mitochondrial Fo complex subunit B1 |
| Database Link: | <u>NP_001679</u> <u>Entrez Gene 515 Human</u> <u>P24539</u> |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| | ATP5PB Goat Polyclonal Antibody – TA329080 |
|-----------------|--|
| Background: | Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) -containing the extramembraneous catalytic core, and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F(0) domain and the peripheric stalk, which acts as a stator to hold the catalytic alpha(3)beta(3) subcomplex and subunit a/ATP6 static relative to the rotary elements. |
| Synonyms: | PIG47 |
| Protein Pathway | s: Alzheimer's disease, Huntington's disease, Metabolic pathways, Oxidative phosphorylation, Parkinson's disease |

Product images:

| | 250kDa 150kDa 100kDa 75kDa | |
|---|-------------------------------------|---|
| | 50kDa 37kDa | TA329080 (0.03ug/ml) staining of Human Heart lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence. |
| - | 25kDa | |
| | 20kDa | |
| | 15kDa | |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US