

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA329365

ESE1 (ELF3) Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-Elf3 antibody: synthetic peptide directed towards the middle region of human Elf3. Synthetic peptide located within the following region: GSCGAGAPSPGSSDVSTAGTGASRSSHSSDSGGSDVDLDPTDGKLFPSDG
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. Note that this product is shipped as lyophilized powder to China customers.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	41 kDa
Gene Name:	E74 like ETS transcription factor 3
Database Link:	<u>NP_004424</u> <u>Entrez Gene 1999 Human</u> <u>P78545</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	ESE1 (ELF3) Rabbit Polyclonal Antibody – TA329365
Background:	ELF3 is a novel, highly tissue-restricted member of the ets transcription factor/oncogene family. ELF3 contains two putative DNA binding domains: an ETS domain, which is unique in that the 5' half shows relatively weak homology to known ets factors, and an A/T hook domain, found in HMG proteins and various other nuclear factors. In contrast to any known ets factors, ELF3 is expressed exclusively in epithelial cells. ELF3 is expected to be a critical regulator of epithelial cell differentiation. ETS factor ELF3 is a transcriptional regulator of angiopoietin-1 gene regulation in the setting of inflammation. ELF3 overexpressed early during human breast tumorigenesis. ELF3 expression confers a transformed and in vitro metastatic phenotype to otherwise normal MCF-12A cells. ELF3 mediates the expression of TGF-beta RII, and the transcriptional inhibition of ets-related transcription factor could be a one of the mechanisms of colonic carcinogenesis. ELF3 is positively and negatively modulated by other interacting proteins including Ku70, Ku86, p300, and CBP.
Synonyms:	EPR-1; ERT; ESE-1; ESX
Note:	lmmunogen sequence homology: Human: 100%; Rabbit: 100%; Pig: 93%; Horse: 92%; Bovine: 86%; Dog: 79%
Protein Families	: Transcription Factors

Product images:



WB Suggested Anti-Elf3 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1:62500; Positive Control: MCF7 cell lysateElf3 is supported by BioGPS gene expression data to be expressed in MCF7

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US