

Product datasheet for TA330486

MURF1 (TRIM63) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

Host: Rabbit

Isotype: IgG
Clonality: Polyclonal

Immunogen: The immunogen for anti-TRIM63 antibody: synthetic peptide directed towards the middle

region of human TRIM63. Synthetic peptide located within the following region:

EQLDKSTKLVETAIQSLDEPGGATFLLTAKQLIKSIVEASKGCQLGKTEQ

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 40 kDa

Gene Name: tripartite motif containing 63

Database Link: NP 115977

Entrez Gene 84676 Human

Q969Q1

Background: This gene encodes a member of the RING zinc finger protein family found in striated muscle

and iris. The product of this gene is localized to the Z-line and M-line lattices of myofibrils, where titin's N-terminal and C-terminal regions respectively bind to the sarcomere. In vitro binding studies have shown that this protein also binds directly to titin near the region of titin containing kinase activity. Another member of this protein family binds to microtubules. Since these family members can form heterodimers, this suggests that these proteins may serve as

a link between titin kinase and microtubule-dependent signal pathways in muscle.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: IRF; MURF1; MURF2; RNF28; SMRZ

Note: Immunogen sequence homology: Dog: 100%; Pig: 100%; Rat: 100%; Horse: 100%; Human:

100%; Mouse: 100%; Bovine: 100%; Rabbit: 100%; Guinea pig: 100%

Product images:



WB Suggested Anti-TRIM63 Antibody Titration: 0.2-1 ug/ml; Positive Control: THP-1 cell lysate