

## **Product datasheet for TA330610**

## **TAF5L Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications:IHC, WBRecommended Dilution:WB, IHCReactivity:HumanHost:RabbitIsotype:IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-TAF5L antibody: synthetic peptide directed towards the N terminal of

human TAF5L. Synthetic peptide located within the following region: LTDSDSQHSHEVMPLLYPLFVYLHLNLVQNSPKSTVESFYSRFHGMFLQN

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 66 kDa

**Gene Name:** TATA-box binding protein associated factor 5 like

Database Link: NP 001020418

Entrez Gene 27097 Human

<u>075529</u>



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. TAF5L encodes a protein that is a component of the PCAF histone acetylase complex and structurally similar to one of the histone-like TAFs, TAF5. The PCAF histone acetylase complex, which is composed of more than 20 polypeptides some of which are TAFs, is required for myogenic transcription and differentiation.

Synonyms: PAF65B


**Note:** Dog: 100%; Pig: 100%; Horse: 100%; Human: 100%; Rabbit: 100%; Guinea pig: 100%; Rat: 92%;

Bovine: 92%; Mouse: 86%

**Protein Families:** Transcription Factors

**Protein Pathways:** Basal transcription factors


## **Product images:**



WB Suggested Anti-TAF5L Antibody Titration: 2.5ug/ml; Positive Control: Jurkat cell lysateTAF5L is supported by BioGPS gene expression data to

be expressed in Jurkat





Rabbit Anti-TAF5L Antibody; Paraffin Embedded Tissue: Human alveolar cell; Cellular Data: Epithelial cells of renal tubule; Antibody Concentration: 4.0-8.0 ug/ml; Magnification: 400X