

Product datasheet for **TA331083**

UGT1A5 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-UGT1A5 antibody: synthetic peptide directed towards the N terminal of human UGT1A5. Synthetic peptide located within the following region: EENFFTLTTYAISWTQDEFDRLLLGHTQSFFETEHELLMKFSRRMAIMNNM
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	57 kDa
Gene Name:	UDP glucuronosyltransferase family 1 member A5
Database Link:	NP_061951 Entrez Gene 54579 Human P35504



[View online »](#)

Background:

This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are considered pseudogenes. Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter.

Synonyms:

UDPGT; UDPGT 1-5; UGT1E

Note:


Human: 100%; Rat: 83%

Protein Families:

Transmembrane

Protein Pathways:

Androgen and estrogen metabolism, Ascorbate and aldarate metabolism, Drug metabolism - cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Pentose and glucuronate interconversions, Porphyrin and chlorophyll metabolism, Retinol metabolism, Starch and sucrose metabolism

Product images:

WB Suggested Anti-UGT1A5 Antibody Titration:
0.2-1 ug/ml; Positive Control: HT1080 cell lysate