

## Product datasheet for **TA332078**

### ABCD4 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for Anti-ABCD4 Antibody: synthetic peptide directed towards the C terminal of human ABCD4. Synthetic peptide located within the following region: FGPHGVLFLPQKPFFTDGLREQVIYPLKEVYPDSGSADDERILRFLELA
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	55 kDa
Gene Name:	ATP binding cassette subfamily D member 4
Database Link:	<a href="#">NP_005041</a> <a href="#">Entrez Gene 5826 Human</a> <a href="#">O14678</a>



[View online »](#)

**Background:** ABCD4 is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ALD subfamily, which is involved in peroxisomal import of fatty acids and/or fatty acyl-CoAs in the organelle. All known peroxisomal ABC transporters are half transporters which require a partner half transporter molecule to form a functional homodimeric or heterodimeric transporter. The function of this peroxisomal membrane protein is unknown. However, it is speculated that it may function as a heterodimer for another peroxisomal ABC transporter and, therefore, may modify the adrenoleukodystrophy phenotype. It may also play a role in the process of peroxisome biogenesis.


**Synonyms:** ABC41; EST352188; MAHCJ; P70R; P79R; PMP69; PXMP1L

**Note:** Immunogen sequence homology: Dog: 100%; Pig: 100%; Rat: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Bovine: 100%; Rabbit: 100%; Guinea pig: 100%; Yeast: 79%

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** ABC transporters

### Product images:



WB Suggested Anti-ABCD4 Antibody Titration: 1.25ug/ml; Positive Control: HepG2 cell lysate; ABCD4 is supported by BioGPS gene expression data to be expressed in HepG2