

Product datasheet for TA332508S

ZAP70 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB 1:500 - 1:2000 Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human ZAP70

Formulation: Store at -20°C (regular) and -80°C (long term). Avoid freeze / thaw cycles. Buffer: PBS with

0.02% sodium azide, 50% glycerol, pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size:

Gene Name: zeta chain of T cell receptor associated protein kinase 70kDa

Database Link: NP 001070

Entrez Gene 22637 MouseEntrez Gene 301348 RatEntrez Gene 7535 Human

P43403

Background: The Syk family protein tyrosine kinase Zap-70 is expressed in T and NK cells and plays a

> critical role in mediating T cell activation in response to T cell receptor (TCR) engagement (1). Following TCR engagement, Zap-70 is rapidly phosphorylated on several tyrosine residues through autophosphorylation and transphosphorylation by the Src family tyrosine kinase Lck

(2-6). Tyrosine phosphorylation correlates with increased Zap-70 kinase activity and downstream signaling events. Expression of Zap-70 is correlated with disease progression

and survival in patients with chronic lymphocytic leukemia (7,8).

Synonyms: SRK; STCD; STD; TZK; ZAP-70



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Natural killer cell mediated cytotoxicity, Primary immunodeficiency, T cell receptor signaling

pathway

Product images:



Western blot analysis of extracts of various cell lines, using ZAP70 antibody.