

Product datasheet for TA332565S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Triosephosphate isomerase (TPI1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:1000

IF/ICC,1:50 - 1:200

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Formulation: Store at -20°C (regular) and -80°C (long term). Avoid freeze / thaw cycles. Buffer: PBS with

0.02% sodium azide, 50% glycerol, pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 27kDa

Gene Name: triosephosphate isomerase 1

Database Link: NP 000356

Entrez Gene 21991 MouseEntrez Gene 24849 RatEntrez Gene 7167 Human

P60174

Background: This gene encodes an enzyme, consisting of two identical proteins, which catalyzes the

isomerization of glyceraldehydes 3-phosphate (G3P) and dihydroxy-acetone phosphate (DHAP) in glycolysis and gluconeogenesis. Mutations in this gene are associated with

triosephosphate isomerase deficiency. Pseudogenes have been identified on chromosomes 1,

4, 6 and 7. Alternative splicing results in multiple transcript variants.

Synonyms: HEL-S-49; TIM; TPI; TPID






Protein Pathways:

Fructose and mannose metabolism, Glycolysis / Gluconeogenesis, Inositol phosphate metabolism, Metabolic pathways

Product images:



Western blot analysis of various lysates using TPI1 Rabbit pAb ([TA332565]) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.