

## **Product datasheet for TA332705S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **IRAKM (IRAK3) Rabbit Polyclonal Antibody**

#### **Product data:**

Product Type: Primary Antibodies

Applications: ELISA, ICC/IF, WB

Recommended Dilution: WB.1:500 - 1:2000

IF/ICC,1:50 - 1:200

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Formulation: Store at -20°C (regular) and -80°C (long term). Avoid freeze / thaw cycles. Buffer: PBS with

0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 68kDa

**Gene Name:** interleukin 1 receptor associated kinase 3

Database Link: NP 009130

Entrez Gene 73914 MouseEntrez Gene 314870 RatEntrez Gene 11213 Human

Q9Y616

**Background:** This gene encodes a member of the interleukin-1 receptor-associated kinase protein family.

Members of this family are essential components of the Toll/IL-R immune signal transduction pathways. This protein is primarily expressed in monocytes and macrophages and functions as a negative regulator of Toll-like receptor signaling. Mutations in this gene are associated with a susceptibility to asthma. Alternate splicing results in multiple transcript variants.

Synonyms: ASRT5; IRAKM






**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Apoptosis, Neurotrophin signaling pathway

# **Product images:**



Western blot analysis of various lysates using IRAK3 Rabbit pAb ([TA332705]) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.