

## Product datasheet for **TA332735S**

### Plakophilin 2 (PKP2) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	WB, 1:500 - 1:2000 ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Formulation:	Store at -20°C (regular) and -80°C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	97kDa
Gene Name:	plakophilin 2
Database Link:	<a href="#">NP_001005242</a> <a href="#">Entrez Gene 67451 Mouse</a> <a href="#">Entrez Gene 287925 Rat</a> <a href="#">Entrez Gene 5318 Human</a> <a href="#">Q99959</a>



[View online »](#)

**Background:**


This gene encodes a member of the arm-repeat (armadillo) and plakophilin gene families. Plakophilin proteins contain numerous armadillo repeats, localize to cell desmosomes and nuclei, and participate in linking cadherins to intermediate filaments in the cytoskeleton. This gene may regulate the signaling activity of beta-catenin and is required to maintain transcription of genes that control intracellular calcium cycling including ryanodine receptor 2, ankyrin-B, triadin, and calcium channel, voltage-dependent, L type, alpha 1C. Mutations in this gene are associated with different inherited cardiac conditions including Arrhythmogenic Cardiomyopathy, Brugada Syndrome, and Idiopathic Ventricular Fibrillation. A processed pseudogene with high similarity to this gene has been mapped to chromosome 12p13.

**Synonyms:**

ARVD9

**Protein Pathways:**

Arrhythmogenic right ventricular cardiomyopathy (ARVC)

**Product images:**


Western blot analysis of various lysates using PKP2 Rabbit pAb (TA332735S) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.