

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA332767

PSME2 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB 1:500 - 1:2000
Reactivity:	Human, Mouse
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Recombinant protein of human PSME2
Formulation:	Store at -20°C (regular) and -80°C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	239
Gene Name:	proteasome activator subunit 2
Database Link:	<u>NP_002809</u> <u>Entrez Gene 19188 MouseEntrez Gene 5721 Human</u> <u>Q9UL46</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


PSME2 Rabbit Polyclonal Antibody – TA332767

Background:The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure
composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4
rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are
composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6
ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase
subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration
and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An
essential function of a modified proteasome, the immunoproteasome, is the processing of
class I MHC peptides. The immunoproteasome contains an alternate regulator, referred to as
the 11S regulator or PA28, that replaces the 19S regulator. Three subunits (alpha, beta and
gamma) of the 11S regulator have been identified. This gene encodes the beta subunit of the
11S regulator, one of the two 11S subunits that is induced by gamma-interferon. Three beta
and three alpha subunits combine to form a heterohexameric ring. Six pseudogenes have
been identified on chromosomes 4, 5, 8, 10 and 13.

Synonyms: PA28B; PA28beta; REGbeta

Protein Pathways: Antigen processing and presentation, Proteasome

Product images:



Western blot analysis of extracts of various cell lines, using PSME2 antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US