

Product datasheet for TA332876

CRYGD Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: WB 1:500 - 1:2000 Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human CRYGD

Formulation: Store at -20°C (regular) and -80°C (long term). Avoid freeze / thaw cycles. Buffer: PBS with

0.02% sodium azide, 50% glycerol, pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 174

Gene Name: crystallin gamma D

Database Link: NP 008822

Entrez Gene 12967 MouseEntrez Gene 24278 RatEntrez Gene 1421 Human

P07320



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are further divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Gamma-crystallins are a homogeneous group of highly symmetrical, monomeric proteins typically lacking connecting peptides and terminal extensions. They are differentially regulated after early development. Four gamma-crystallin genes (gamma-A through gamma-D) and three pseudogenes (gamma-E, gamma-F, gamma-G) are tandemly organized in a genomic segment as a gene cluster. Whether due to aging or mutations in specific genes, gamma-crystallins have been involved in cataract formation.

Synonyms: CACA; CCA3; CCP; cry-g-D; CRYG4; CTRCT4; PCC

Protein Families: Druggable Genome

Product images:



Western blot analysis of extracts of various cell lines, using CRYGD antibody.