

Product datasheet for TA333196

EIF4EBP1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB 1:500-1:2000

Reactivity: Human, Mouse, Rat

Modifications: Phospho-specific

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A phospho specific peptide corresponding to residues surrounding T36 of human EIF4EBP1

Formulation: Store at -20°C (regular) and -80°C (long term). Avoid freeze / thaw cycles. Buffer: PBS with

0.02% sodium azide, 50% glycerol, pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 118

Gene Name: eukaryotic translation initiation factor 4E binding protein 1

Database Link: NP 004086

Entrez Gene 13685 MouseEntrez Gene 116636 RatEntrez Gene 1978 Human

Q13541

Background: This gene encodes one member of a family of translation repressor proteins. The protein

directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of

mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of

mRNA translation.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: 4E-BP1; 4EBP1; BP-1; PHAS-I

Protein Pathways: Acute myeloid leukemia, ErbB signaling pathway, Insulin signaling pathway, mTOR signaling

pathway

Product images:



Western blot analysis of extracts from MDA-MB-435 cells, using Phospho-eIF4EBP1-T36 antibody (TA333196). | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per

lane. | Blocking buffer: 3% BSA.