

Product datasheet for TA333303

p73 (TP73) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB 1:500 - 1:2000

Reactivity: Human

Modifications: Phospho-specific

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A phospho specific peptide corresponding to residues surrounding Y99 of human TP73

Formulation: Store at -20°C (regular) and -80°C (long term). Avoid freeze / thaw cycles. Buffer: PBS with

0.02% sodium azide, 50% glycerol, pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 636

Gene Name: tumor protein p73

Database Link: NP 005418

Entrez Gene 7161 Human

O15350



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes a member of the p53 family of transcription factors involved in cellular responses to stress and development. It maps to a region on chromosome 1p36 that is frequently deleted in neuroblastoma and other tumors, and thought to contain multiple tumor suppressor genes. The demonstration that this gene is monoallelically expressed (likely from the maternal allele), supports the notion that it is a candidate gene for neuroblastoma. Many transcript variants resulting from alternative splicing and/or use of alternate promoters have been found for this gene, but the biological validity and the full-length nature of some variants have not been determined.

Synonyms: P73

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Neurotrophin signaling pathway, p53 signaling pathway

Product images:



Western blot analysis of extracts from 293 cells, using Phospho-TP73-Y99 antibody (TA333303). | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA.