

## **Product datasheet for TA333726**

## SLC4A5 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Immunogen:** The immunogen for Anti-SLC4A5 Antibody: synthetic peptide directed towards the middle

region of human SLC4A5. Synthetic peptide located within the following region:

SIAHIDSLKMETETSAPGEQPQFLGVREQRVTGIIVFILTGISVFLAPIL

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Purification:** Affinity Purified

Conjugation: Unconjugated

**Store** at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 126 kDa

**Gene Name:** solute carrier family 4 member 5

Database Link: NP 067019

Entrez Gene 57835 Human

Q9BY07

**Background:** This gene encodes a member of the sodium bicarbonate cotransporter (NBC) family, part of

the bicarbonate transporter superfamily. Sodium bicarbonate cotransporters are involved in intracellular pH regulation and electroneural or electrogenic sodium bicarbonate transport. This protein is thought to be an integral membrane protein. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some

variants has not been determined. [provided by RefSeg, Jul 2008]



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: NBC4; NBCe2

**Note:** Immunogen sequence homology: Dog: 100%; Pig: 100%; Rat: 100%; Horse: 100%; Human:

100%; Mouse: 100%; Bovine: 100%; Rabbit: 100%; Zebrafish: 100%; Guinea pig: 100%

**Protein Families:** Transmembrane

## **Product images:**



WB Suggested Anti-SLC4A5 Antibody Titration: 0.2-1 ug/ml; Positive Control: THP-1 cell lysate