

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA334022

#### **TEF Rabbit Polyclonal Antibody**

### **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	The immunogen for Anti-TEF Antibody: synthetic peptide directed towards the middle region of human TEF. Synthetic peptide located within the following region: RKDEGRKEAAEGKEQGLAPLVVQTDSASPLGAGHLPGLAFSSHLHGQQFF
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. Note that this product is shipped as lyophilized powder to China customers.
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	33 kDa
Gene Name:	TEF, PAR bZIP transcription factor
Database Link:	<u>NP_003207</u> <u>Entrez Gene 7008 Human</u> <u>Q10587</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


## **TEF Rabbit Polyclonal Antibody – TA334022**

Background:	TEF (thyrotroph embryonic factor) is a member of the PAR bZip (proline and acidic amino acid-rich basic leucine zipper) transcription factor family. It accumulates with robust circadian rhythms in tissues with high amplitudes of clock gene expression. Thyrotroph embryonic factor (TEF), a transcription factor, is a member of the PAR (proline and acidic amino acid-rich) subfamily of basic region/leucine zipper (bZIP) transcription factors. It is expressed in a broad range of cells and tissues in adult animals, however, during embryonic development, TEF expression appears to be restricted to the developing anterior pituitary gland, coincident with the appearance of thyroid-stimulating hormone, beta (TSHB). Indeed, TEF can bind to, and transactivate the TSHB promoter. It shows homology (in the functional domains) with other members of the PAR-bZIP subfamily of transcription factors, which include albumin D boxbinding protein (DBP), human hepatic leukemia factor (HLF) and chicken vitellogenin genebinding protein (VBP); VBP is considered the chicken homologue of TEF. Different members of the subfamily form heterodimers, and share DNA-binding, and transcriptional regulatory properties.
Synonyms:	KIAA1655
Note:	Immunogen sequence homology: Dog: 100%; Horse: 100%; Human: 100%; Rat: 92%; Mouse:

lmmunogen sequence homology: Dog: 100%; Horse: 100%; Human: 100%; Rat: 92%; Mouse: 92%; Sheep: 92%; Bovine: 92%; Rabbit: 85%

Protein Families: Transcription Factors

# **Product images:**



WB Suggested Anti-TEF Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1:12500; Positive Control: 293T cell lysate

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US