

Product datasheet for TA334246

TBX4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-TBX4 antibody: synthetic peptide directed towards the middle region

of human TBX4. Synthetic peptide located within the following region: LRVARLQSKEYPVISKSIMRQRLISPQLSATPDVGPLLGTHQALQHYQHE

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 60 kDa

Gene Name: T-box 4

Database Link: NP 060958

Entrez Gene 9496 Human

P57082



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

TBX4 Rabbit Polyclonal Antibody - TA334246

Background: The TBX4 gene is a member of a phylogenetically conserved family of genes that share a

common DNA-binding domain, the T-box. T-box genes encode transcription factors involved in the regulation of developmental processes. This gene is the human homolog of mouse Tbx4, which is closely linked to Tbx2 on mouse chromosome 11. Similarly this gene, like TBX2, maps to human chromosome 17. Expression studies in mouse and chicken show that Tbx4 is expressed in developing hindlimb, but not in forelimb buds, suggesting a role for this gene in

regulating limb development and specification of limb identity.

Synonyms: ICPPS; SPS

Note: Immunogen Sequence Homology: Dog: 100%; Pig: 100%; Rat: 100%; Goat: 100%; Horse: 100%;

Human: 100%; Mouse: 100%; Sheep: 100%; Bovine: 100%; Rabbit: 100%; Guinea pig: 100%;

Zebrafish: 79%

Product images:



WB Suggested Anti-TBX4 Antibody Titration: 0.2-1 ug/ml; Positive Control: HepG2 cell lysate