

# **Product datasheet for TA335787**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SP2 transcription factor (SP2) Rabbit Polyclonal Antibody

### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB

Reactivity: Human

Host: Rabbit

**Isotype:** IgG

Clonality: Polyclonal

Immunogen: The immunogen for Anti-SP2 Antibody: synthetic peptide directed towards the N terminal of

human SP2. Synthetic peptide located within the following region: SDPQTSMAATAAVSPSDYLQPAASTTQDSQPSPLALLAATCSKIGPPAVE

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Purification:** Affinity Purified

Conjugation: Unconjugated

**Store** at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 65 kDa

**Gene Name:** Sp2 transcription factor

Database Link: NP 003101

Entrez Gene 6668 Human

Q02086





Background:

Sp family proteins are sequence-specific DNA-binding proteins characterized by an aminoterminal trans-activation domain and three carboxy-terminal zinc finger motifs. This protein contains the least conserved DNA-binding domain within the Sp subfamily of proteins, and its DNA sequence specificity differs from the other Sp proteins. It localizes primarily within subnuclear foci associated with the nuclear matrix, and can activate or in some cases repress expression from different promoters. This gene encodes a member of the Sp subfamily of Sp/XKLF transcription factors. Sp family proteins are sequence-specific DNA-binding proteins characterized by an amino-terminal trans-activation domain and three carboxy-terminal zinc finger motifs. This protein contains the least conserved DNA-binding domain within the Sp subfamily of proteins, and its DNA sequence specificity differs from the other Sp proteins. It localizes primarily within subnuclear foci associated with the nuclear matrix, and can activate or in some cases repress expression from different promoters. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

**Synonyms:** KIAA0048; Sp2 transcription factor

Note: Immunogen Sequence Homology: Pig: 100%; Human: 100%; Bovine: 100%; Rabbit: 100%;

Mouse: 93%; Horse: 88%; Dog: 86%; Rat: 86%; Guinea pig: 86%

**Protein Families:** Transcription Factors

### **Product images:**



WB Suggested Anti-SP2 Antibody Titration: 0.2-1 ug/ml; ELISA Titer: 1: 62500; Positive Control: Transfected 293T