

## **Product datasheet for TA336342**

## rioduct datasileet for TASSOS4

## **SIRT7 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: Western Blot: 1-3 ug/ml

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

**Immunogen:** A mixture of synthetic peptides corresponding to amino acids 35-51 and 361-377 of human

SIRT7 was used as immunogen.

Formulation: PBS containing 0.05% BSA, 0.05% Sodium Azide. Store at 4C short term. Aliquot and store at -

20C long term. Avoid freeze-thaw cycles.

**Concentration:** lot specific

**Purification:** Protein G purified

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Gene Name: sirtuin 7

Database Link: NP 057622

Entrez Gene 51547 Human

Q9NRC8

**Background:** SIRT7 is a human member of a family of proteins called Sirtuins (Sir2-like proteins) and are

present in prokaryotes and eukaryotes. All Sir2-like proteins have a sirtuin core domain, which contains a series of sequence motifs conserved in organisms ranging from bacteria to humans. Bacterial, yeast and mammalian sirtuins are able to metabolize NAD and possibly at as mono-ADP-ribosyltransferases. The enzymatic function of sirtuins is not yet completely understood but recent reports of histone-activated Sir2-mediated NAD metabolism and NAD-

activated Sir2-mediated histone deacetylation suggest a possible coupled reciprocal

activation mechanism involving interactions of Sir2 with NAD and the N epsilon-acetyl-lysine

groups of acetylated histones.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: SIR2L7

**Protein Families:** Druggable Genome, Transcription Factors

## **Product images:**



Western Blot: SIRT7 Antibody TA336342 - Analysis of SIRT7 in A) human liver and B) PBMC lysate using this antibody at 1 ug/ml.