

Product datasheet for **TA336398**

IKK gamma (IKBKG) Mouse Monoclonal Antibody [Clone ID: 72C627]

Product data:

| | |
|-----------------------|--|
| Product Type: | Primary Antibodies |
| Clone Name: | 72C627 |
| Applications: | Simple Western, WB |
| Recommended Dilution: | Western Blot: 2-5 ug/ml, Simple Western: 1:200 |
| Reactivity: | Human, Mouse |
| Host: | Mouse |
| Isotype: | IgG1 |
| Clonality: | Monoclonal |
| Immunogen: | This antibody was raised against a His-tagged full-length human IKK3/IKKg protein. |
| Formulation: | PBS containing 0.05% BSA, 0.05% Sodium Azide. Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. |
| Concentration: | lot specific |
| Purification: | Protein G purified |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Gene Name: | inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase gamma |
| Database Link: | NP_001093326 Entrez Gene 16151 Mouse Entrez Gene 8517 Human Q9Y6K9 |



[View online »](#)

Background:

NF- κ B (nuclear factor κ B) is sequestered in the cytoplasm by I κ B family of inhibitory proteins that mask the nuclear localization signal of NF- κ B thereby preventing translocation of NF- κ B to the nucleus. External stimuli such as tumor necrosis factor or other cytokines results in phosphorylation and degradation of I κ B releasing NF- κ B dimers. NF- κ B dimer subsequently translocates to the nucleus and activates target genes. Synthesis of I κ Ba is autoregulated. I κ B proteins are phosphorylated by I κ B kinase complex consisting of at least three proteins, IKK1/a, IKK2/b, and IKK3/g. IKK3/g preferentially interacts with IKK2/b and is required for activation of IKK complex. IKK3/g is also known as NEMO (NF- κ B Essential MODulator). Recent data suggest that the human T-cell leukemia virus type I Tax oncoprotein that activates NF- κ B binds neither to IKKa nor IKKb, but complexes directly with IKKg. This suggests that IKKg may be a key molecule acting as an adapter for onco-protein specific signaling to IKKa and IKKb.

Synonyms:


AMCBX1; FIP-3; FIP3; Fip3p; IKK-gamma; IKKAP1; IKKG; IMD33; IP; IP1; IP2; IPD2; NEMO; ZC2HC9

Protein Families:


Druggable Genome, Transcription Factors

Protein Pathways:

Acute myeloid leukemia, Adipocytokine signaling pathway, Apoptosis, B cell receptor signaling pathway, Chemokine signaling pathway, Chronic myeloid leukemia, Cytosolic DNA-sensing pathway, Epithelial cell signaling in Helicobacter pylori infection, MAPK signaling pathway, NOD-like receptor signaling pathway, Pancreatic cancer, Pathways in cancer, Primary immunodeficiency, Prostate cancer, RIG-I-like receptor signaling pathway, Small cell lung cancer, T cell receptor signaling pathway, Toll-like receptor signaling pathway

Product images:

Simple Western: IKK gamma Antibody (72C627) TA336398 - Simple Western lane view shows a specific band for IKK gamma in 0.05 mg/ml of Jurkat lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Western Blot: IKK gamma Antibody (72C627) TA336398 - Analysis using IKK gamma (NEMO) antibody. Lysate from 1) human Jurkat and 2) mouse NIH 3T3 cells probed with IKK gamma antibody at 2 ug/ml.