

Product datasheet for **TA336612**

GAP43 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	WB: 1:10000, IF: 1:1000, IHC: 1-2 ug/ml, IHC-F: 1-2 ug/ml, IHC-P: 1-2 ug/ml
Reactivity:	Human, Mouse, Rat, Chicken, Primate
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	C-terminal peptide of rat and mouse GAP43, which is KEDPEADQEHA, with an N-terminal Cys added to allow chemical coupling to KLH carrier protein.
Formulation:	PBS, 30% glycerol, 0.1% Sodium Azide. Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Concentration:	lot specific
Purification:	Immunogen affinity purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	43 kDa
Gene Name:	growth associated protein 43
Database Link:	NP_002036 Entrez Gene 14432 Mouse Entrez Gene 29423 Rat Entrez Gene 2596 Human P17677



[View online »](#)

Background:

GAP43 (growth associated protein 43, also called B-50, PP46, calmodulin-binding protein P-57, neuromodulin, neuron growth-associated protein 43, protein F1) is a major component of motile "growth cones" that form the tips of elongating axons as well as the synaptic plasma membranes. GAP43 is associated with nerve growth process and plays a key role in axonal as well as dendritic filopodia induction. It has been detected in a complex containing FGFR4, NCAM1, CDH2, PLCG1, FRS2, SRC, SHC1, GAP43 and CTTN, and binds calmodulin with superior affinity in the absence of Ca²⁺ than in its presence. Intracellular Ca²⁺ regulates PKC mediated phosphorylation of GAP43 and affects GAP43's ability to bind calmodulin. GAP43 is integral to growth cone formation, neurite outgrowth, and the development of a functional cerebral cortex. Aberrant expression of GAP43 has been seen in patients diagnosed with schizophrenia and Alzheimer's disease. Three GAPs (Growth associated proteins) were discovered, and the number 43 comes from the apparent SDS-PAGE molecular weight of the one named GAP43. The HGNC name for this protein is, not surprisingly, GAP43. Later work showed that GAP43 does not run on SDS-PAGE in a fashion which accurately reflects its molecular weight, and that GAP43 proteins from different species may run at different apparent molecular weights. Partly due to these features GAP43 was independently discovered by several different groups and therefore has several alternate names, such as protein F1, pp46, neuromodulin, neural phosphoprotein B-50 and calmodulin-binding protein P-57. In each case the number reflects the apparent SDS-PAGE molecular weight, and underlines the unusual SDS-PAGE mobility properties of this molecule. Mammalian GAP43 proteins contains only 226-243 amino acids, and so the real molecular weight is 23.61-25.14 kDa (to perform such calculations yourself see this link). GAP43 is one of many highly negatively charged extended molecules which lack well defined tertiary structure and contain few hydrophobic residues and which run anomalously on SDS-PAGE. Other examples are CAP23, MARCKS, microtubule associated proteins MAP2, tau and the Neurofilament subunits. GAP43 has been extensively studied and is known to be a major protein kinase C substrate and to bind calmodulin avidly. GAP43 is anchored to the plasma membrane by palmitoylation modifications.

Synonyms:


B-50; PP46

Note:


This GAP43 antibody is useful for Immunocytochemistry/Immunofluorescence, Immunohistochemistry on paraffin-embedded and frozen sections and Western blot, where it recognizes a band at 43 kDa.

Product images:


Western Blot: GAP43 Antibody TA336612 - Western blots of homogenate of cow cerebellum stained with RPCA-GAP43. A prominent band running at ~43kDa represents the full length GAP43.




Immunohistochemistry: GAP43 Antibody TA336612 - Region CA1-CA4 of the hippocampus showing strong GAP-43 staining in cell processes and neuropil using TA336612.




Immunohistochemistry-Paraffin: GAP43 Antibody TA336612 - Review image from confirmed customer on mouse E15.5 paraffin sections.



Immunocytochemistry/Immunofluorescence: GAP43 Antibody TA336612 - Immunofluorescence of GAP-43 (green), a molecular marker of neurite outgrowth, demonstrates intense staining in overexpressing wild-type PS-1 (E) PC-12 cells. (Teo, et al, 2005)



Immunocytochemistry/Immunofluorescence: GAP43 Antibody TA336612 - Mixed neuron-glia cultures stained with RPCA-GAP43 (red), blue is DNA staining.



Immunocytochemistry/Immunofluorescence:
GAP43 Antibody TA336612 - Rat E18 mixed
neuron/glia cultures with rabbit GAP43 (red) and
5B10, mouse monoclonal to MAP-tau (green).