

Product datasheet for TA336631

Product data:

Product Type: Primary Antibodies

GAPDH Rabbit Polyclonal Antibody

Applications: WB

Recommended Dilution: WB: 1:2000-1:10000

Reactivity: Human, Mouse, Xenopus, Yeast

Host: Rabbit

Clonality: Polyclonal

Immunogen: The epitope recognized by this antibody maps to a region between residues 200 and 250 of

human GAPDH using the numbering given in entry NP_002037.2 (GeneID 2597).

Formulation: Tris-citrate/phosphate, pH 7, 0.1% Sodium azide. Store at 4C. Do not freeze.

Concentration: lot specific

Purification: Immunogen affinity purified

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 38 kDa

Gene Name: glyceraldehyde-3-phosphate dehydrogenase

Database Link: NP 002037

Entrez Gene 14433 MouseEntrez Gene 2597 Human

P04406



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

GAPDH (glyceraldehyde-3-phosphate dehydrogenase or GAPD) is a key enzyme in glycolytic pathway, wherein it catalyzes the first step by converting D-glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate. It localizes mainly in the cytoplasm from where it translocates to nucleus following S-nitrosylation and interaction with SIAH1. Nuclear GAPDH implicates in transcription, RNA transport, DNA replication and apoptosis, via its nitrosylase activity which mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2, PRKDC etc. GAPDH also regulates the organization/assembly of cytoskeleton and facilitates CHP1-dependent microtubule - membrane associations. It is a component of GAIT (gamma interferon-activated inhibitor of translation) complex which mediates IFN-gamma-induced transcript-selective translation inhibition in inflammation. Because of its expression as housekeeping protein in most cell types, GAPDH is often used as a control molecule in various genes expression studies, however, recent evidence has shown the association of its altered expression with neurodegenerative pathologies such as Huntington disease, Alzheimer's disease etc., and elevated GAPDH mRNA/protein expression levels have been seen in pancreatic, lung and prostate cancers.

Synonyms: G3PD; GAPD; HEL-S-162eP


Note: This GAPDH antibody is useful for Western blot, where a band is observed ~36 kDa. The

antibody does not work in Immunoprecipitation.

Protein Families: ES Cell Differentiation/IPS


Protein Pathways: Alzheimer's disease, Glycolysis / Gluconeogenesis, Metabolic pathways

Product images:




Western Blot: GAPDH Antibody TA336631 - Detection of Human and Mouse GAPDH Whole cell lysate from HeLa (5, 15 and 50 mcg) and mouse NIH3T3 (M; 50 mcg). Another Affinity purified rabbit anti-GAPDH antibody used at 0.04 mcg/ml. Chemiluminescence with an





Western Blot: GAPDH Antibody TA336631 - Western blot analysis of GAPDH (TA336631) using NIH3T3 whole cell lysate [NB800-PC8].



Western Blot: GAPDH Antibody TA336631 - Western blot analysis of GAPDH (TA336631) using whole normal mouse brain lysate [NB820-59657].