

## Product datasheet for **TA336645**

### SCARB1 Rabbit Polyclonal Antibody

#### Product data:

|                         |                                                                                                                                                                                                                                                                                |
|-------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Type:           | Primary Antibodies                                                                                                                                                                                                                                                             |
| Applications:           | Block/Neutralize, FC, ICC/IF, IHC, IP, WB                                                                                                                                                                                                                                      |
| Recommended Dilution:   | Immunohistochemistry-Paraffin, Immunohistochemistry-Frozen (Negative): 5 ug/ml, Knockout Validated, Flow Cytometry: 1:400, Block/Neutralize, Immunohistochemistry: 1:10-1:500, Immunocytochemistry/ Immunofluorescence: 1:50-1:1000, Immunoprecipitation, Western Blot: 1:1000 |
| Reactivity:             | Human, Mouse, Rat                                                                                                                                                                                                                                                              |
| Host:                   | Rabbit                                                                                                                                                                                                                                                                         |
| Clonality:              | Polyclonal                                                                                                                                                                                                                                                                     |
| Immunogen:              | A peptide from the extracellular domain (residues 230-380) of Scavenger Receptor-BI/BII that was expressed as two tandem copies in bacteria using the pET system.                                                                                                              |
| Formulation:            | Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.                                                                                                                                                                                                                   |
| Concentration:          | lot specific                                                                                                                                                                                                                                                                   |
| Purification:           | Whole antisera                                                                                                                                                                                                                                                                 |
| Conjugation:            | Unconjugated                                                                                                                                                                                                                                                                   |
| Storage:                | Store at -20°C as received.                                                                                                                                                                                                                                                    |
| Stability:              | Stable for 12 months from date of receipt.                                                                                                                                                                                                                                     |
| Predicted Protein Size: | 82 kDa                                                                                                                                                                                                                                                                         |
| Gene Name:              | scavenger receptor class B member 1                                                                                                                                                                                                                                            |
| Database Link:          | <a href="#">NP_005496</a><br><a href="#">Entrez Gene 20778 Mouse</a> <a href="#">Entrez Gene 25073 Rat</a> <a href="#">Entrez Gene 949 Human</a><br><a href="#">Q8WTV0</a>                                                                                                     |



[View online »](#)

**Background:**

SR-BI (scavenger receptor class BI, SCARB1) is a member of CD36 receptor family and act as a major receptor for HDL for facilitating selective lipid uptake from lipoproteins in several tissues. SR-BI gene gives rise to at least two mRNA splice variants (SR-BI and SR-BII), and interestingly, SR-BII variant is the main isoform in terms of mRNA levels in certain tissues e.g. SR-BII mRNA is 48-fold higher than SR-BI in brain cells; and protein levels reaching 10-15% of SR-BI in the liver. Similar differential SR-BII protein expression levels have also been detected in considerable amounts in other tissues, such as testes, retinal pigment epithelial cells, and spleen. SR-BII isoform differs from SR-BI only in its entirely different, yet highly conserved, cytoplasmic C-terminus which has been proposed to contains a signal responsible for marked alterations in cellular distribution as well as cellular trafficking of SR-BII compared with SR-BI. In contrast to SR-BI, SR-BII mediates the rapid uptake of significant amounts of HDL into endosomal recycling compartment by a pathway which is different from selective lipid uptake at the cell surface (mediated by SR-BI). Because SR-BII influence cellular cholesterol trafficking/homeostasis in a manner distinct from SR-BI, the relative expression as well as functional activities of these two isoforms create a potential means of regulating selective lipid transfer between HDL and cells.

**Synonyms:**


CD36L1; CLA-1; CLA1; HDLQTL6; SR-BI; SRB1

**Note:**


This SR-BI/SR-BII antibody is useful for Flow Cytometry (PMID: 22097902), functional studies (customer feedback), Immunocytochemistry, Western blot and Immunohistochemistry. In Western blot a band is observed at ~ 82 kDa in tissues that express SR-BI and/or SR-BII such as liver, ovary, adrenal glands, and to as lesser extent testes, heart and mammary glands.

**Protein Families:**


Druggable Genome, Transmembrane

**Product images:**



Immunocytochemistry/Immunofluorescence: SR-BI/SR-BII Antibody TA336645 - SR-BI/SR-BII antibody was tested in HeLa cells with DyLight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and DyLight 550 (red).



Flow Cytometry: SR-BI/SR-BII Antibody TA336645 - SR-BI/SR-BII antibody was tested at 1:400 in NIH-3T3 cells using an Alexa Fluor 488 secondary (shown in purple). M1 is defined by unstained cells.



Immunohistochemistry-Paraffin: SR-BI/SR-BII Antibody TA336645 - IHC analysis of formalin-fixed paraffin-embedded tissue section of normal human liver using 5 ug/ml concentration of SR-BI/SR-BII antibody. Specific and expected granular membrane-cytoplasmic staining was observed in the hepatocytes. [40X Magnification]



Western Blot: SR-BI/SR-BII Antibody TA336645 - Detection of RED-1 in 80 ug of total mouse liver lysates. Lane 1: wild-type mice, Lane 2: SR-BI deficient mice.