

Product datasheet for TA336710

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NFH (NEFH) Mouse Monoclonal Antibody [Clone ID: 9B12]

Product data:

Product Type: Primary Antibodies

Clone Name: 9B12
Applications: IF, WB

Recommended Dilution: WB: 1:10000, IF: 1:1000

Reactivity: Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Native Neurofilament Heavy protein purified from bovine spinal cord

Formulation: Preservative: 0.05% Sodium Azide. Aliquot and store at -20C or -80C. Avoid freeze-thaw

cycles.

Concentration: lot specific

Purification: Ascites

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 200 kDa

Gene Name: neurofilament, heavy polypeptide

Database Link: NP 066554

Entrez Gene 24587 Rat

P12036





Background:

Neuronal intermediate filaments (NIF) are found specifically on neurons where they play key role in maintaining neuronal morphology as well as in regenerating myelinated axons. NIF contain three subunits: a light polypeptide (NEFL/NFL), a medium polypeptide (NEFM/NFM), and a heavy polypeptide (NEFH/NFH), with molecular weights of 68, 160, and 212 kD respectively. 200kD neurofilament heavy (NEFH) is one of the major components of neuronal cytoskeleton NIF and NEFH gene was originally proposed as a DNA marker for presymptomatic diagnosis in neurofibromatosis type 2 (NF2). NEHF facilitate interactions between NIF and brain mitochondria with binding efficiency dependent upon mitochondrial membranes potential. NEFH and beta-catenin are constituents in postsynaptic density (PSD) in CNS and NEFH is essential for maintaining normal cell integrity. Diminished NEFH levels are seen in certain cancers including human autonomic nerve tumors or central neurocytomas, prostate cancer, clear-cell epithelioid tumor and small cell lung carcinoma. Fibroblasts overexpressing NEFH are distorted, multinucleated, and possess inclusions, typical changes seen in neuropathies, and defects in NEFH are linked to susceptibility to ALS.

Synonyms: NFH


Note: This 200kDa Neurofilament Heavy antibody is useful for

Immunocytochemistry/Immunofluorescence and Western Blot.

Protein Families: Druggable Genome


Protein Pathways: Amyotrophic lateral sclerosis (ALS)

Product images:



Analysis of different tissue lysates using mouse mAb to NF-H, TA336710, dilution 1:10000 (Green): [1] protein standard, [2] rat spinal cord [3] mouse spinal cord, and [4] cow spinal cord. Strong band at about 200-220kDa corresponds to the major phosphorylated form of the NF-H subunit. Smaller proteolytic fragments of NF-H are also detected in some preparations.





Immunocytochemistry/Immunofluorescence: 200kDa Neurofilament Heavy Antibody (9B12) TA336710 - Mixed neuron/glial cultures stained with TA336710 (red) and a rabbit GFAP antibody (green). Axonal profiles are stained in red, while astrocytic cells are revealed in green. Nuclei are revealed with a fluorescent DNA stain (blue).