

Product datasheet for TA336726

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PDK4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: Immunocytochemistry/ Immunofluorescence: 1:500, Western Blot: 2 ug/ml

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide made to an internal portion of the human PDK4 protein (within residues

200-250). [Swiss-Prot# Q16654]

Formulation: PBS, 0.05% Sodium Azide. Store at 4C short term. Aliquot and store at -20C long term. Avoid

freeze-thaw cycles.

Concentration: lot specific

Purification: Immunogen affinity purified

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 49 kDa

Gene Name: pyruvate dehydrogenase kinase 4

Database Link: NP 002603

Entrez Gene 5166 Human

Q16654



Background:

PDK4 (pyruvate dehydrogenase kinase isoform 4) is expressed ubiquitously in the mitochondrion matrix of tissues with highest expression seen in liver, heart and skeletal muscles. PDK4 is regulated by glucocorticoids, retinoic acid and insulin, and is found elevated with diabetes, fasting and other conditions where fatty acids act as energy source. Under starvation/diabetic conditions, enhanced glucocorticoids levels activate glucocorticoid receptor thereby recruiting p300/CBP coactivator complex to PDK4 gene where it cooperatively interact with FOXO factors and activated RARs to make a stable transcriptional initiation complex responsible for PDK4 activation. Insulin promotes inactivation and translocation of FOXO factors from nucleus to cytoplasm and thereby suppresses glucocorticoid-stimulated PDK4 gene transcription. PDK4 inhibits the mitochondrial pyruvate dehydrogenase complex (PDC) via its E1 alpha subunit's phosphorylation, thus contributing to glucose metabolism regulation. Insulin resistance is associated with PDC dysregulation in skeletal muscles and excessive insulin on the other hand downregulates PDK4 expression. PDK4 exerts conservation of glucose as well as three-carbon compounds such as lactate, alanine, pyruvate etc. which serve as substrates for gluconeogenesis and this conservation helps maintain euglycemia during starvation, however, exacerbates hyperglycemia in diabetes.

Synonyms: FLJ40832


Note: This PDK4 antibody is useful for Immunocytochemistry/Immunofluorescence and Western

blot, where a band is seen at ~49 kDa. In ICC/IF mitochondrial staining was observed in HeLa

cells.


Protein Families: Druggable Genome, Protein Kinase

Product images:



Immunocytochemistry/Immunofluorescence: PDK4 Antibody TA336726 - PDK4 antibody was tested in HeLa cells with Dylight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and Dylight 550 (red).





Western Blot: PDK4 Antibody TA336726 - Western Blot on human heart protein lysate.