

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA336787

TIGAR Rabbit Polyclonal Antibody

Product data:

| Product Type: | Primary Antibodies |
|-----------------------|---|
| Applications: | IHC, WB |
| Recommended Dilution: | Immunohistochemistry: 1:10-1:500, Western Blot: 1:5000, Immunohistochemistry-Paraffin: 1:10-1:500 |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Immunogen: | A genomic peptide made to an internal region of the human TIGAR protein (within residues 50-200). [Swiss-Prot Q9NQ88] |
| Formulation: | PBS, 0.05% Sodium Azide. Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. |
| Concentration: | lot specific |
| Purification: | Immunogen affinity purified |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Gene Name: | TP53 induced glycolysis regulatory phosphatase |
| Database Link: | <u>NP_065108</u> <u>Entrez Gene 319801 MouseEntrez Gene 57103 Human</u> <u>Q9NQ88</u> |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

TIGAR Rabbit Polyclonal Antibody - TA336787

Background: TIGAR (TP53-induced glycolysis and apoptosis regulator), a member of phosphoglycerate mutase family, is a target gene of p53 and it regulates glucose metabolism in cancer cells by acting as a fructose-bisphosphatase hydrolyzing fructose-2,6-bisphosphate (Fru-2,6-P2) and Fru-1,6-P2 leading to decreased Fru-2,6-P2 levels with subsequent reduction in PFK1 (phosphofructosekinase-1) activity as well as negative regulation of glycolysis. TIGAR also reduces cellular ROS through its translocation to the mitochondria where it interacts with HK2 leading to enhanced HK2 activity, regulation of mitochondrial membrane potential, and decreased mitochondrial ROS which ultimately protects the cells from DNA damage-induced apoptosis. Moreover, TIGAR mediated negative regulation of glycolysis leads to redirection of glycolytic metabolic intermediates towards pentose phosphate pathway's oxidative branch that follows increased cellular NADPH production, ROS scavenging by GSH and thus a lower sensitivity of cells to oxidative stress-associated apoptosis, including that induced by p53. Because of TIGAR's anti-apoptotic/protective role, it has been proposed to implicate in cellcycle arrest and DNA repair effector response in cells that suffer from repairable dose of genotoxic/cellular insult.

Synonyms:

Note:

C12orf5; FR2BP

This TIGAR antibody is useful for Immunohistochemistry-paraffin sections and Western blot. In Western blot bands are seen ~30 kDa, representing TIGAR, and ~47 kDa. At this time we cannot explain the band at 47 kDa. This antibody does not appear to work in ICC.

Product images:



Western Blot: TIGAR/C12orf5 Antibody TA336787 - Analysis of TIGAR in HeLa whole cell extracts.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemistry: TIGAR/C12orf5 Antibody TA336787 - Analysis of TIGAR in mouse muscle.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US