

Product datasheet for TA336868

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SUV420h1 (KMT5B) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, IHC, WB

Recommended Dilution: Immunohistochemistry: 1:200, Immunocytochemistry/ Immunofluorescence: 1:100-1:300,

Immunohistochemistry-Paraffin: 1:200, Western Blot (Negative)

Reactivity: Human, Mouse

Host: Rabbit

Clonality: Polyclonal

Immunogen: A synthetic peptide made to an internal portion of the human SUV420h1 protein (between

residues 100-150) [UniProt Q4FZB7]

Formulation: PBS with 30% Glycerol, 0.05% Sodium Azide. Store at 4C short term. Aliquot and store at -20C

long term. Avoid freeze-thaw cycles.

Concentration: lot specific

Purification: Affinity purified Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: lysine methyltransferase 5B

Database Link: NP 060105

Entrez Gene 225888 MouseEntrez Gene 51111 Human

Q4FZB7

Background: SUV420H1 (Histone-lysine N-methyltransferase 5B, KMT5B) is a histone methyltransferase

that specifically di- and trimethlyates lysine 20 of H4 (H4 K20Me3), an epigenetic mark of transcriptional repression. This mark, along with trimethylation of histone 3 lysine 9 (H3 K9me3) and monomethylation of H3 lysine 27 (H3 K27me1), define genomic regions of constitutive heterochromatin that are poor in activated genes. This enzyme is essential for development, with Suv420h1 knockout mice proving lethal perinatally and showing overall growth retardation. The RB1 family of proteins have been shown to interact with, and target,

KMT5B to histones, enabling the enzyme to function.





SUV420h1 (KMT5B) Rabbit Polyclonal Antibody - TA336868

Synonyms: CGI-85; CGI85; SUV420H1


Note: This SUV420h1 antibody is useful for Immunocytochemistry/Immunofluorescence and IHC-

paraffin embedded sections. In ICC/IF, nuclear staining was observed in A431 cells. In IHC-P, staining was observed in the nucleus of mouse epidermis tissue. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended. This


antibody is not recommended for Western Blot.

Protein Families: Druggable Genome
Protein Pathways: Lysine degradation

Product images:



Immunocytochemistry/Immunofluorescence: SUV420h1 Antibody TA336868 - SUV420h1 antibody was tested in A431 cells with FITC (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and Dylight 550 (red).



Immunohistochemistry: SUV420h1 Antibody TA336868 - IHC analysis of SUV420h1 in mouse epidermis using DAB with hematoxylin counterstain.